

Flat Spacetime Vacuum in Loop Quantum Gravity

A. Miković *

Departamento de Matemática e Ciências de Computação, Universidade
Lusófona, Av. do Campo Grande, 376, 1749-024, Lisboa, Portugal

Abstract

We construct a state in the loop quantum gravity theory with zero cosmological constant, which should correspond to the flat spacetime vacuum solution. This is done by defining the loop transform coefficients of a flat connection wavefunction in the holomorphic representation which satisfies all the constraints of quantum General Relativity and it is peaked around the flat space triads. The loop transform coefficients are defined as spin foam state sum invariants of the spin networks embedded in the spatial manifold for the $SU(2)$ quantum group. We also obtain an expression for the vacuum wavefunction in the triad representation, by defining the corresponding spin networks functional integrals as $SU(2)$ quantum group state sums.

*E-mail address: amikovic@ulusofona.pt

1 Introduction

In [1] it was demonstrated that the invariants of the embedded spin networks for the $SU(2)$ BF theory can be interpreted as the loop transform coefficients of a flat connection wavefunction which satisfies all the constraints of quantum General Relativity (GR) in the Ashtekar formulation. Since the flat spacetime can be represented as a flat connection solution with the flat metric constraint, this opened up a possibility of constructing a flat metric vacuum state in the framework of loop quantum gravity [2, 3, 4].

The approach of constructing a vacuum state by using the spin network invariants was initiated by Smolin for the case of non-zero cosmological constant, see [5] for a review and references. In this case, a solution of the quantum constraints has been already known, which is the Kodama state [6]. The corresponding loop transform coefficients are given by the $SU(2, \mathbf{C})$ Chern-Simons (CS) theory invariants of the embedded spin networks, and it was conjectured that these invariants are given by an analytic continuation of the $SU_q(2)$ quantum group spin networks evaluations at a root of unity [5]. The quantum group evaluations of the spin networks are given by the Kauffman brackets [7, 8, 9]. Consequently, one has to use the quantum spin networks, i.e. the spin networks where the $SU(2)$ irreducible representations (irreps) are restricted by $j \leq k/2$, where $k \in \mathbf{N}$ and $k = 6\pi/\Lambda l_P^4$, where Λ is the cosmological constant and l_P is the Planck length. In the $\Lambda = 0$ case, the CS theory is replaced by the BF theory, and in order to define the corresponding spin network invariants one must use the same category of the $SU_q(2)$ irreps [1], but then k is an arbitrary natural number.

The state constructed in [1] corresponds to a flat connection wavefunction which is not peaked around any particular value of the triads, and hence cannot be a good description of the flat metric vacuum. The same problem also appears in the case of non-zero cosmological constant, because the Kodama state is not peaked around any particular value of the triads[†].

In this paper we extend the construction of [1] to the case when the flat connection wavefunction is peaked around the flat metric values of the triads, so that we obtain a state in the spin network basis which could be considered as a flat space vacuum. In section 2 we give a review and some basic formulas of the holomorphic representation. These formulas are then used in section 3

[†]Classically, self/anti-self dual connection solution describes de-Sitter/anti-de-Sitter solution only if the spatial metric is flat.

to construct a flat connection wavefunction which is peaked around the flat space triads. In section 4 the corresponding spin network basis coefficients are defined as the $SU_q(2)$ spin foam state sum invariants. Since the vacuum wavefunction satisfies the quantum constraints in the holomorphic representation, in section 5 we construct the wavefunction in the triad representation by defining the corresponding functional integrals as $SU_q(2)$ state sums. In section 6 we present our conclusions, and in the Appendix we present the calculation of the expansion coefficients for the gauge invariant plane waves.

2 Holomorphic representation

The Ashtekar formulation of GR [10] is based on the complex canonical variables

$$A_i^a = -i\omega_i^a(e) + \tilde{p}_i^a \quad , \quad E_a^i = \tilde{e}_a^i \quad , \quad (1)$$

where $\tilde{e}_a^i = \sqrt{h}e_a^i$, e 's are the inverse triads, h is the determinant of the metric on the three-manifold Σ , $\omega(e)$ is the torsion free spin connection and $\tilde{p}$ is the canonically conjugate variable to $\tilde{e}$ [‡]. The constraints of GR become polynomial in terms of the new variables, and the Hamiltonian constraint is given by

$$C_H = \epsilon_{abc} \left(E^{ia} E^{jb} F_{ij}^c + \Lambda \epsilon_{ijk} E^{ia} E^{jb} E^{kc} \right) \quad , \quad (2)$$

where Λ is the cosmological constant.

The polynomiality of the constraints simplifies the quantization of the theory; however, one has to find a representation of the complex canonical variables which is consistent with the reality conditions, i.e.

$$A^\dagger - A = 2i\omega(E) \quad , \quad E^\dagger = E \quad . \quad (3)$$

In the case of finitely many canonical variables, this is resolved via the holomorphic representation [11]. Given a Heisenberg algebra $[p, q] = i$, one can introduce creation and annihilation operators

$$a = p + i\Omega'(q) \quad , \quad a^\dagger = p - i\Omega'(q) \quad , \quad (4)$$

[‡]Here we use the conventions such that the spin connection is the imaginary part of the complex connection A . Hence the curvature two-form is given by $F^a = dA^a + i\epsilon_{abc}A^b \wedge A^c$ where the $SU(2)$ Lie algebra generators T_a satisfy $[T_a, T_b] = i\epsilon_{abc}T_c$. An $SU(2)$ group element is given by $g = e^{i\Theta^a T_a}$, where Θ^a are real parameters.

where Ω is a given function of q and $\Omega' = \frac{d\Omega}{dq}$. Since the operators a and $a^\dagger$ are not hermitian, their eigenvalues α and $\bar{\alpha}$ will be complex numbers, where $\bar{\alpha}$ is the complex conjugate of α . Let us for the sake of convinience denote the eigenvector of $a^\dagger$ as $|\alpha\rangle$, so that

$$a^\dagger|\alpha\rangle = \bar{\alpha}|\alpha\rangle \quad . \quad (5)$$

These eigenstates can be expressed in the q representation as

$$\langle q|\alpha\rangle = e^{-i\bar{\alpha}q + \Omega(q)} \quad , \quad (6)$$

so that for an arbitrary state $|\Psi\rangle$ we can define $\Psi(\alpha)$ as

$$\Psi(\alpha) = \langle \alpha|\Psi\rangle = \int dq \langle \alpha|q\rangle \langle q|\Psi\rangle = \int dq e^{i\alpha q + \Omega(q)} \Phi(q) \quad . \quad (7)$$

The complex variable function $\Psi(\alpha)$ will be holomorphic, and one can define the holomorphic representation as

$$a|\Psi\rangle \rightarrow \alpha\Psi(\alpha) \quad , \quad q|\Psi\rangle \rightarrow i\Psi'(\alpha) \quad . \quad (8)$$

Given a holomorphic wavefunction $\Psi(\alpha)$, one can go back to the q representation via the contour integral

$$\Phi(q) = e^{-\Omega(q)} \int_{C=\mathbf{R}} d\alpha e^{-i\alpha q} \Psi(\alpha) \quad , \quad (9)$$

so that in order to obtain the usual wavefunction we need to know $\Psi(\alpha)$ for the real values of α .

In the case of GR one can define the holomorphic representation as [11]

$$\hat{A}|\Psi\rangle \rightarrow A\Psi[A] \quad , \quad \hat{E}|\Psi\rangle \rightarrow i\frac{\delta\Psi}{\delta A} \quad , \quad (10)$$

where

$$\Psi[A] = \langle A|\Psi\rangle = \int \mathcal{D}E \langle A|E\rangle \langle E|\Psi\rangle = \int \mathcal{D}E e^{i\int_\Sigma d^3x AE + \Omega[E]} \Phi[E] \quad , \quad (11)$$

and $\frac{\delta\Omega}{\delta E} = \omega(E)$. Given a $\Psi[A]$ functional, we can obtain the $\Phi[E]$ functional as

$$\Phi[E] = e^{-\Omega[E]} \int_{A\in\mathbf{R}} \mathcal{D}A e^{-i\int_\Sigma d^3x AE} \Psi[A] \quad . \quad (12)$$

Although the expresions in the GR case are a straightforward generalization of the finite-dimensional ones, the key difference is that all the integrals that are used in the finite-dimensional case become functional integrals, and these have to be defined. For our purposes it will suffice to define the functional integral (12).

3 Flat-connection wavefunctions

In [1] it was pointed out that the wavefunctions $\delta(F)\Psi_0(A)$ are solutions of the $\Lambda = 0$ GR constraints in the holomorphic representation. Since at the classical level the $F = 0$ solutions include the flat metric solution, i.e. when $A = i\omega(E)$ then $F(A) = iR_3$, where R_3 is the scalar curvature of Σ . One can then use the $\delta(F)\Psi_0(A)$ wavefunctions to construct a flat metric vacuum state [1].

Let $|0\rangle$ be a state corresponding to $\delta(F)\Psi_0(A)$, then one can represent this state in the basis of diffeo invariant spin network states $|\gamma\rangle$ as [3]

$$|0\rangle = \sum_{\gamma} \langle\gamma|0\rangle |\gamma\rangle \quad , \quad (13)$$

where

$$\langle\gamma|0\rangle = \int \mathcal{D}A W_{\gamma}[A] \delta(F)\Psi_0[A] = I_{\gamma} \quad . \quad (14)$$

The complex number I_{γ} is a topological invariant for a spin network γ embedded into the three-manifold Σ , and the formal functional integral expression (14) can be used to define this invariant. This can be done by replacing the Σ by a simplicial complex corresponding to a triangulation of Σ and then by using the dual one-complex to define the integral over the connections as an integral over the group elements associated to the dual edge holonomies. This finite-dimensional integral can be then regulated via the quantum $SU(2)$ group at a root of unity, which can be represented as a spin foam state sum. This was done in [1] for the case of $\Psi_0 = 1$.

Note that in the expression (14) the integration is over the complex A , so that one has to use the $SU(2, \mathbf{C})$ group [§]. Since the relevant category of irreps is always over $\mathbf{C}$, and we do not require unitarity of the quantum group irreps, then the categories of finite-dimensional irreps of $SU_q(2, \mathbf{R})$ and $SU_q(2, \mathbf{C})$ are equivalent.

It was also argued in [1] that the state $\Psi(A) = \delta(F)$ can not correspond to a flat metric vacuum because it does not favor any particular value of the operator $\hat{E}$. It was then suggested to take a nontrivial $\Psi_0(A)$, which would correspond to a state which is sharply peaked around the flat metric value E_0 of $\hat{E}$. Observe that if $\Psi_0(A)$ is an eigenstate of $\hat{E}$, then $\Psi(A) = \delta(F)\Psi_0(A)$

[§]The Lie algebra $su(2, \mathbf{C})$ is isomorphic to the Lie algebra $sl(2, \mathbf{C})$, when $sl(2, \mathbf{C})$ is considered as a complex vector space. If we consider $sl(2, \mathbf{C})$ as a real vector space, then it is isomorphic to $su(2) \oplus su(2)$.

is not, but we can still consider the state $\Psi(A)$ as a state which is sharply peaked around E_0 . The reason is that the formula (14) can be rewritten as

$$\langle \gamma | 0 \rangle = \int \mathcal{D}A^* W_\gamma[A^*] \Psi_0[A^*] \quad , \quad (15)$$

i.e. as a functional integral over the flat connections A^* , see also [12]. Hence one can reinterpret $|0\rangle$ as a state from a vector space corresponding to a quantization of the space of flat connections. Let E^* be the canonical conjugate of $Re A^*$, then the holomorphic representation is defined by the same formulas as in the case of a non-flat connection A . In particular we have

$$\begin{aligned} \Psi_0[A^*] &= \langle A^* | \Psi_0 \rangle = \int \mathcal{D}E^* \langle A^* | E^* \rangle \langle E^* | \Psi_0 \rangle \\ &= \int \mathcal{D}E^* e^{i \int_\Sigma d^3x A^* E^* + \Omega[E^*]} \Phi[E^*] \quad , \end{aligned} \quad (16)$$

so that if we choose for Φ an eigenfunctional of $\hat{E}^*$ for the flat-metric eigenvalue E_0 , which is given formally by $\delta(E^* - E_0)$, we then obtain

$$\Psi_0[A^*] = e^{i \int_\Sigma d^3x A^* E_0 - \Omega[E_0]} = e^{i \int_\Sigma d^3x A^* E_0} \quad , \quad (17)$$

since $\Omega(E_0) = 0$.

Therefore this analysis suggests to take for the flat metric vacuum a state $|0\rangle$ whose coefficients in the spin network basis are given by the formal expression

$$\langle \gamma | 0 \rangle = \int \mathcal{D}A W_\gamma[A] \delta(F) e^{i \int_\Sigma d^3x A E_0} \quad . \quad (18)$$

In the next section we will define this functional integral as a spin-foam state sum for the quantum $SU(2)$ group at a root of unity.

4 Flat metric spin network invariants

Let $T(\Sigma)$ be a simplicial complex corresponding to a triangulation of Σ . We consider only the triangulations where the dual one-complex of $T(\Sigma)$ is a four-valent graph Γ , see Fig. 1. We can define (18) by taking

$$\mathcal{D}A = \prod_l dg_l \quad , \quad (19)$$

where l labels the edges of Γ , g_l is an $SU(2)$ group element corresponding to the edge holonomy and dg_l is the usual group measure (Haar measure). Then $\delta(F)$ can be defined as

$$\prod_f \delta(g_f) \quad , \quad (20)$$

where f labels the faces of Γ and $g_f = \prod_{l \in \partial f} g_l$. The group delta function is defined as

$$\delta(g) = \sum_j (2j+1) \text{Tr} D^{(j)}(g) \quad , \quad (21)$$

where j is the spin and $D^{(j)}$ is the corresponding representation matrix.

We will embed the spin network γ into the graph Γ by identifying a subset of vertices V' of Γ with the vertices of γ and then we will assign to the edges of Γ which connect the vertices from V' the irreps of the spin network γ (one can have more than one irrep of γ associated to the same edge of Γ). Then

$$W_\gamma[A] = \langle \prod_{l \in L'} \prod_{j_l \in J_\gamma} D^{(j_l)}(g_l) \prod_{v \in V'} C^{(\iota_v)} \rangle \quad , \quad (22)$$

where L' is a set of edges of Γ labelled by the irreps of γ , J_γ is the set of irreps of γ , $C^{(\iota)}$ are the intertwiner tensors for the intertwiners of γ and $\langle \rangle$ denotes the $SU(2)$ trace [13].

As far as $\Psi_0 = e^{i \int_\Sigma d^3x \langle A E_0 \rangle}$ is concerned, it can be discretized as

$$\Psi_0 = \exp i \sum_{l \in L} \langle A_l E_\Delta^0 \rangle = \prod_{l \in L} e^{i \langle A_l E_\Delta^0 \rangle} \quad , \quad (23)$$

where

$$E_\Delta^a = \int_\Delta E^{ai} \epsilon_{ijk} dx^j \wedge dx^k \quad , \quad (24)$$

and Δ is a triangle dual to the edge l . Due to the Peter-Weyl theorem, one has

$$e^{i A^a E_a} = \sum_j (c_j(E))_\beta^\alpha D_\alpha^{(j)\beta}(e^{i A^a T_a}) \quad , \quad (25)$$

where

$$(c_j(E))_\beta^\alpha = (2j+1) \int_{SU(2)} dg \bar{D}_\alpha^{(j)\beta}(g) f(g) \quad , \quad f(g) = f(e^{i A^a T_a}) = e^{i A^a E_a} \quad . \quad (26)$$

We then get

$$\Psi_0 = \sum_{j_1, \dots, j_L} c_{\beta_1}^{\alpha_1} \cdots c_{\beta_L}^{\alpha_L} D_{\alpha_1}^{(j_1)\beta_1}(g_1) \cdots D_{\alpha_L}^{(j_L)\beta_L}(g_L) \quad . \quad (27)$$

Note that in order to have a good definition of I_γ , the function Ψ_0 should be gauge invariant. This translates into the invariance under $g_l \rightarrow h_v^{-1} g_l h_{v'}$, where v and v' are the ends of l and h 's are arbitrary group elements. This invariance requires that one finds an approximation

$$c_{\beta_1}^{\alpha_1} \cdots c_{\beta_L}^{\alpha_L} \approx \sum_{\iota} f(E, \iota) C_{\beta_1 \dots \beta_L}^{\alpha_1 \dots \alpha_L(\iota)} \quad , \quad (28)$$

where $C^{(\iota)}$ are the intertwiner tensors[¶]. The simplest way to achieve this is to replace the plane-wave expansion (25) by a gauge-invariant expression

$$(e^{iAE})_{inv} = \sum_j C_j(E) Tr D^{(j)}(e^{iAT}) \quad , \quad (29)$$

where $(2j+1)C_j(E) = Tr(c_j(E))$. Then by taking the trace of both sides of the approximation (28), and assuming that all the f 's are equal, one obtains

$$f(E, \iota) = \frac{C_{j_1}(E_1) \cdots C_{j_L}(E_L)}{(2j_1+1) \cdots (2j_L+1)} \quad . \quad (30)$$

It is not difficult to calculate the coefficients C_j , see the Appendix. An interesting feature is that they are non-zero only for $j \leq \epsilon = |E|/l_P^2$. The C_j 's are given by the expression

$$C_j(E) = [I_0(j) - I_0(j+1) + I_2(j) - I_2(j+1)] \quad , \quad (31)$$

where

$$I_0(j) = \frac{\theta(\epsilon - j)}{\sqrt{\epsilon^2 - j^2}} \quad , \quad I_2(j) = \frac{1 - 2j^2/\epsilon^2}{\sqrt{\epsilon^2 - j^2}} \theta(\epsilon - j) \quad , \quad (32)$$

where θ is the step function ($\theta(x) = 0$ for $x \leq 0$ and $\theta(x) = 1$ for $x > 0$).

In the case when $(E_0)_i^a = \delta_i^a$ (flat space), it is natural to take $\epsilon = 1$, so that the expansion (29) terminates at $j = 1$. The C_1 coefficient is then divergent, and one would have to decide how to regularize it. The corresponding I_γ will be given by the integral

$$I_\gamma = \sum_{j'_i, \iota'} \int \prod_l dg_l \tilde{C}_{j'_l}(E_0) \prod_f \delta(g_f) W_\gamma(g_{\nu}, j, \iota) \langle C^{(\iota')} D^{(j'_1)}(g_1) \cdots D^{(j'_L)}(g_L) \rangle \quad , \quad (33)$$

[¶]One cannot have an equality because the coefficients $c_\alpha^\beta(E)$ are not the group tensors.

where $(2j+1)\tilde{C}_j = C_j$. This integral can be represented by a state sum

$$I_\gamma = \sum_{j_f, \iota_l, j'_l, \iota'_v} \prod_f (2j_f + 1) \prod_l \tilde{C}_{j'_l}(E_0) \prod_v A_v(j_f, \iota_l; j'_l, \iota'_v; j, \iota) \quad . \quad (34)$$

The vertex amplitudes A are evaluations of the vertex spin networks associated to the spin foam $\{\Gamma, j_f, \iota_l\}$ carrying the spin networks $\{\Gamma, j'_l, \iota'_v\}$ and $\{\gamma, j', \iota'\}$. The vertex spin networks can be obtained in the following way: draw the circuit diagram Γ_c for Γ (each circuit is labelled by a j_f) and then draw the $\Gamma(j', \iota')$ and the $\gamma(j, \iota)$ spin networks on the Γ_c diagram. This gives the circuit diagram $\Gamma_c(\Gamma, \gamma)$, see Fig. 2. Then shrink each edge of $\Gamma_c(\Gamma, \gamma)$ to a point, and label these points with the intertwiners ι_l . These rules follow from the graphical representation of the group integrations [14], see Fig. 3. In this way one obtains the vertex spin networks which are the tetrahedral spin networks with the additional four-vertices j', ι' and additional lines and vertices corresponding to the spin network γ , see Fig. 4.

The state sum for I_γ is regularized by passing to the category of finite dimensional irreps of $SU_q(2)$ where q is a root of unity. Let $q = e^{2\pi i/(k+2)}$, $k \in \mathbf{N}$, then the $SU(2)$ irreps satisfy $j \leq k/2$ and

$$I_\gamma = \sum_{j_f, \iota_l, j'_l, \iota'_v} \prod_f \dim_q(j_f) \prod_l C_{j'_l}(E_0) \prod_v A_v^{(q)}(j_f, \iota_l; j'_l, \iota'_v; j, \iota) \quad , \quad (35)$$

where $\dim_q$ is the quantum dimension and $A_v^{(q)}$ is the quantum group evaluation of the vertex spin network. These spin networks evaluations can be calculated by using the formulas from [9].

5 The triad representation

Given the invariants I_γ for the flat-space vacuum, one can construct the corresponding wavefunction as

$$\Psi[A] = \sum_\gamma I_\gamma \bar{W}_\gamma[A] \quad . \quad (36)$$

Since A is complex, we want to obtain the usual wavefunction, i.e. the function of the real argument. We then use the formula (12) to go to the triad representation, so that

$$\Phi[E] = \sum_\gamma I_\gamma \bar{\Phi}_\gamma[E] \quad , \quad (37)$$

where

$$\Phi_\gamma[E] = e^{-\Omega[E]} \int_{A \in \mathbf{R}} \mathcal{D}A e^{-i \int_\Sigma d^3x A E} W_\gamma[A] \quad . \quad (38)$$

This functional integral can be also defined as a state sum via the simplicial decomposition of Σ . As in the previous section we use the formulas (19), (22) and (23) so that

$$\Phi_\gamma[E] = \int \prod_l dg_l \prod_{l \in L} e^{-i A_l E \Delta} \langle \prod_{l \in L'} \prod_{j_l \in J_\gamma} D^{(j_l)}(g_l) \prod_{v \in V'} C^{(\iota_v)} \rangle \quad . \quad (39)$$

By using the gauge invariant approximation (28) we obtain the state sum

$$\Phi_\gamma[E] = \sum_{j', \iota', \tilde{\iota}} C_{j'_1}(E_1) \cdots C_{j'_L}(E_L) \prod_{v \in V'} \mathcal{J}_{\Gamma, \gamma}^{(v)}(j', \iota'; j, \iota; \tilde{\iota}) \quad , \quad (40)$$

where $\mathcal{J}_{\Gamma, \gamma}^{(v)}(j', \iota'; j, \iota; \tilde{\iota})$ are evaluations of the vertex spin networks obtained by a composition of the spin network $\{\Gamma, j', \iota'\}$ and the spin network $\{\gamma, j, \iota\}$.

The vertex spin networks can be obtained by using the following rules: draw the graph of the spin network γ close to the graph of the spin network Γ , such that the resulting graph corresponds to the embedding of the spin network γ into the one-complex Γ , see Fig. 5. Then use the graphical rules for the group integration, see Fig. 3, so that each edge of the $\Gamma \cup \gamma$ graph is cut into two vertices. These new vertices will carry the intertwiners $\tilde{\iota}$, if they are four-valent or higher. If they are two-valent vertices, there will be a factor of the inverse dimension of the corresponding γ irrep. If the two new vertices are one-valent (this happens in the case when an edge of $\Gamma \cup \gamma$ carries only an irrep of the spin network Γ) such vertices are not included, because they will carry the trivial identity irrep. In this way one obtains non-trivial spin networks only for the vertices of Γ which are close to the vertices of γ , see Fig. 6.

The state sum (40) can be then regularized by using the same category of the finite-dimensional irreps of the $SU_q(2)$ quantum group as in the case of the invariant I_γ , i.e. $q = e^{2\pi i/(k+2)}$, $k \in \mathbf{N}$ and $j \leq k/2$.

6 Conclusions

We expect that the state sum (35) for I_γ should be a topological invariant because it was based on the diffeomorphism invariant formal expressions

(18) and (34). This means that I_γ should be independent of the choice of a triangulation of Σ , which means independence of the dual one-complex Γ . Still, one should check the triangulation invariance because it may require different than the usual normalization factors in the spin network amplitudes. This will require checking the invariance under the Pachner moves, see [16] for the case when Σ is a two-dimensional manifold and $\Psi_0 = 1$.

Note that the state sum I_γ corresponds to a new type of a spin foam, due to the presence of a nontrivial function Ψ_0 . Beside labelling the faces of the two-complex Γ with arbitrary group irreps, we also have to label the edges of Γ with arbitrary group irreps.

The expression for the wavefunction in the triad representation (37) will have the form

$$\Phi[E] = \sum_{\gamma} f_{\gamma, \Gamma}(E_1, \dots, E_L) \quad . \quad (41)$$

Since the number of vertices of a spin network γ satisfies $V_\gamma \leq V_\Gamma$, this means that the expression for $\Phi[E]$ will involve a sum over different Γ 's, or equivalently a sum over different triangulations of Σ . Note that summing over triangulations is a way of obtaining diffeomorphism invariant expressions when the individual terms are not diffeomorphism invariant. This raises a possibility to approximate Φ by taking a sufficiently big graph Γ , i.e. a sufficiently fine triangulation of Σ , which would then truncate the infinite sum (41) to a finite sum over the γ 's such that $V_\gamma \leq V_\Gamma$.

Given the functional $\Phi[E]$, the central question is whether it has a good semi-classical limit. This amounts to showing that an appropriately defined semi-classical spacetime metric $g_{\mu\nu}(t, x)$ ($t \in \mathbf{R}$, $x \in \Sigma$) satisfies quantum corrected Einstein equations

$$R_{\mu\nu} + l_P c_1 (\nabla R)_{\mu\nu} + l_P^2 [c_2 (R^2)_{\mu\nu} + c_3 (\nabla^2 R)_{\mu\nu}] + \dots = 0 \quad , \quad (42)$$

where $x^\mu = (t, x)$. The metric $g_{\mu\nu}$ should be an effective classical metric associated to the vacuum state, and a straightforward way to define $g_{\mu\nu}$ is to take that the corresponding spatial metric h_{ij} is given by

$$(\det h) h^{ij}(t, x) = \langle 0 | \hat{E}_a^i(x) \hat{E}^{ja}(x) | 0 \rangle \quad . \quad (43)$$

However, the main problem with this approach is to determine the time evolution parameter t . It will be a function of the parameters which appear in the vacuum state, and it is not obvious how to do this. This is the difficult problem of the time variable in canonical quantum gravity [17], and it is

related to the problem of the choice of the scalar product and the normalizability of the $|0\rangle$ state.

A less straightforward, but more promising approach is to use the De Broglie-Bohm formalism [18]. Then the effective classical metric will be a solution of the quantum equations of motion given by

$$f_i^a(\dot{E}, E, N_\alpha) = \frac{\delta S[E]}{\delta E_a^i(x)} \quad , \quad (44)$$

where $f(\dot{E}, E, N_\alpha)$ is the expression for the $\tilde{p}$ from the canonical formalism, N_α are the lagrange multipliers and $S[E]$ is the phase of the wavefunction $\Phi[E]$. This is a technically simpler approach than the expectation value approach, and the additional advantages are that there is no problem with the interpretation of the wavefunction of the universe^{||} and there is no need for the specification of a scalar product in order to verify the semi-classical limit.

In the case when $\Lambda \neq 0$, the Hamiltonian constraint can be solved in the holomorphic representation via the Kodama wavefunction [6]

$$\Psi_K(A) = \exp \left(\frac{1}{\Lambda} \int_\Sigma \text{Tr}(A \wedge dA + i \frac{2}{3} A \wedge A \wedge A) \right) . \quad (45)$$

One can then construct the corresponding state in the spin network basis via the spin network invariants for the $SU(2)$ CS theory [5]

$$I_\gamma = \int \mathcal{D}A W_\gamma[A] e^{S_{CS}/\Lambda} \quad . \quad (46)$$

When $\Sigma = S^3$ one can argue that this invariant should be defined as a Wick rotation of the Kauffman bracket evaluation for the spin network γ . More precisely, the evaluation for $q = e^{2\pi i/(k+2)}$, where $k \in \mathbf{N}$ and $\Lambda l_P^4 = 6\pi/k$, is analytically continued by $k \rightarrow ik$ [5].

It has been conjectured that the corresponding state in the spin network basis corresponds to the de-Sitter vacuum spacetime when $\Lambda > 0$ [5]. However, the main problem with this conjecture is that the Kodama wavefunction is not peaked around any particular eigenvalue of the $\hat{E}$ operator, since

$$\hat{E}\Psi_K = \frac{1}{\Lambda}\hat{F}\Psi_K \quad . \quad (47)$$

^{||}The wavefunction is interpreted as a wave which guides the point particles, and hence there is no need for the external observer because the physical objects and their properties exist independently of the acts of measurement.

Hence, exactly as in the $\Lambda = 0$ case, Ψ_K cannot be a very good description the de-Sitter vacuum, where the spatial metric is also flat.

One can then try to modify the Kodama wavefunction $\Psi_K(A)$ by multiplying it by a $\Psi_0(A)$ which is peaked around the E_0 eigenvalue, but unlike the $\Lambda = 0$ case, the product $\Psi_K(A)\Psi_0(A)$ does not solve the Hamiltonian constraint. However, if one takes

$$\Psi_0(A) = \delta(F - \Lambda E_0) \quad , \quad (48)$$

then $\Psi_K\Psi_0$ almost solves the Hamiltonian constraint, i.e. $\hat{C}_H\Psi_K\Psi_0 = 0$ when $F \neq \Lambda E_0$, while in the reduced phase space (A^*, E^*) , determined by the constraint $F = \Lambda E_0$, one has that Ψ_K is an eigenstate of the $\hat{E}^*$ operator, see (47). This then suggests to define I_γ via the expression

$$I_\gamma = \int \mathcal{D}A \delta(F - \Lambda E_0) W_\gamma[A] e^{S_{CS}/\Lambda} \quad , \quad (49)$$

where E_0 are the flat space triads. It remains to be seen how useful is this expression for formulating a state sum invariant, but given such an invariant one can obtain the wavefunction in the triad representation by using the formulas of the section 5.

APPENDIX

An $SU(2)$ group element g can be represented as

$$g = e^{i\Theta^a\sigma_a} = \cos(\Theta/2) I + 2in^a\sigma_a \sin(\Theta/2) \quad (50)$$

where σ_a are the Pauli matrices, $n^a = \Theta^a/\Theta$ and $\Theta^2 = \Theta^a\Theta_a$. The trace of the $D^{(j)}(g)$ matrix is given by

$$\chi^{(j)}(g) = \frac{\sin(j + \frac{1}{2})\Theta}{\sin(\Theta/2)} \quad . \quad (51)$$

Since $g_l = e^{iA_l^a\sigma_a}$, then the gauge invariant plane-wave coefficients are given by

$$C_j(E) = \frac{1}{32\pi^2} \int_{\mathbf{R}^3} d^3A \left(\frac{\sin(A/2)}{A/2} \right)^2 \frac{\sin(j + \frac{1}{2})A}{\sin(A/2)} e^{iA^a E_a} \quad , \quad (52)$$

see [14]. This integral can be written as

$$\frac{1}{4\pi} \int_0^\infty dA \int_0^\pi d\theta \sin^2 \theta \sin(A/2) \sin(j + \frac{1}{2})A e^{iAE \cos \theta} \quad , \quad (53)$$

which becomes an integral

$$\frac{1}{4\pi} \int_0^\infty dA [\cos(jA) - \cos(j+1)A] [J_0(AE) + J_2(AE)] \quad , \quad (54)$$

due to the formula

$$J_n(z) = \frac{i^{-n}}{\pi} \int_0^\pi e^{iz \cos \theta} \cos(n\theta) \quad . \quad (55)$$

By using

$$\int_0^\infty J_n(at) \cos(bt) dt = \frac{\cos(n \arcsin(b/a))}{\sqrt{a^2 - b^2}} \quad , \quad 0 \leq b < a \quad (56)$$

and

$$\int_0^\infty J_n(at) \cos(bt) dt = \frac{-a^n \sin(\pi n/2)}{\sqrt{b^2 - a^2} [b + \sqrt{b^2 - a^2}]^n} \quad , \quad 0 < a < b \quad , \quad (57)$$

see [15], one obtains (31).

ACKNOWLEDGEMENTS

I would like to thank Nuno Costa Dias and Joao Prata for the discussions. Work supported by the FCT grants POCTI/FNU/49543/2002 and POCTI/MAT/45306/2002.

References

- [1] A. Miković, Class. Quant. Grav. 20 (2003) 3483
- [2] C. Rovelli and L. Smolin, Nucl. Phys. B 331 (1990) 80
- [3] C. Rovelli and L. Smolin, Phys. Rev. D 52 (1995) 5743
- [4] C. Rovelli, Living Rev. Rel. 1 (1998) 1

- [5] L. Smolin, Quantum gravity with a positive cosmological constant, preprint hep-th/0209079
- [6] H. Kodama, Prog. Theor. Phys. 80 (1988) 1024
- [7] L.H. Kauffman, Int. J. Mod. Phys. A5 (1990) 93
- [8] L.H. Kauffman and S.L. Lins, Temperley-Lieb Recoupling Theory and Invariants of 3-Manifolds, Princeton University Press (1994)
- [9] J.S. Carter, D.E. Flath and M. Saito, The Classical and Quantum 6j-Symbols, Mathematical Notes 43, Princeton University Press (1995)
- [10] A. Ashtekar, Phys. Rev. Lett. 57 (1986) 2244
- [11] H. Kodama, Int. J. Mod. Phys. D 1 (1992) 439
- [12] J. Baez, Knots and quantum gravity: progress and prospects, preprint gr-qc/9410018
- [13] R. Oeckl and H. Pfeiffer, Nucl. Phys. B598 (2001) 400
- [14] L. Friedel and K. Krasnov, Adv. Theor. Math. Phys. 2 (1999) 1183
- [15] M. Abramovitz and I.A. Stegun, Handbook of Mathematical Functions, Dover Publications (1972) New York
- [16] A. Miković, String theory and quantum spin networks, preprint hep-th/0307141
- [17] C. Isham, Canonical quantum gravity and the problem of time, preprint gr-qc/9210011
- [18] P.R. Holland, The Quantum Theory of Motion: An Account of the de Broglie-Bohm Casual Interpretation of Quantum Mechanics, Cambridge University Press (1993)




Figure I: The dual one-complex graph Γ of a triangulation of S^3 with two tetrahedra.




Figure II: The graph $\Gamma_c(\Gamma, \gamma)$ for a spin network $\gamma = \theta_5$ embedded in S^3 whose one-complex Γ is given by that of Fig. 1. The small circles around the edges of Γ_c indicate the group integrations.




Figure III: The graphical representation of the group integrations.




Figure IV: The vertex spin networks for the graph from Fig. 2.




Figure V: The graph $\Gamma \cup \gamma$ for $\Gamma = \theta_4$ and $\gamma = \theta_5$.




Figure VI: The vertex spin networks for the graph from Fig. 5.